

Longest Cycles in Triangle-Free Graphs

MIN AUNG

*Fachbereich Mathematik, Technische Universität Berlin,
Strasse des 17. Juni 135, 1000 Berlin 12, Federal Republic of Germany*

Communicated by J. A. Bondy

Received July 4, 1986

We prove that, for each longest cycle in a 2-connected triangle-free graph G of minimum valence δ on at most $6\delta - 6$ vertices, $G - C$ has at most one edge. It follows that such a graph contains a longest cycle which is also dominating.

© 1989 Academic Press, Inc.

1. INTRODUCTION

We consider only finite simple graphs. Given a graph G and a subgraph K of G , $V(K)$ denotes the vertex set of K , and the subgraph of G induced by $V(G) - V(K)$ is denoted by $G - K$ or by $G - V(K)$. We abbreviate $|K| := |V(K)|$. Cycles and paths in G are considered as subgraphs of G . In particular, $|Q| := |V(Q)|$ for a path Q . A cycle C in G is dominating if $G - C$ has no edge. While considering a graph G and its subgraphs, δ denotes the minimum valence of G .

In this paper we shall prove

THEOREM 1.1. *Let C be a longest cycle in the 2-connected triangle-free graph G . If $|G| \leq 6\delta - 6$, then $G - C$ cannot have two edges.*

In fact, we will separately prove the following two propositions.

PROPOSITION 1.2. *Let C be a longest cycle in the 2-connected triangle-free graph G . If $G - C$ contains a component on at least three vertices, then $|G| \geq 6\delta - 5$.*

PROPOSITION 1.3. *Let C be a longest cycle in the connected triangle-free graph G . If $G - C$ has two distinct components each of which contains exactly two vertices, then $|C| \geq 6\delta - 6$ and $|G| \geq 6\delta - 2$.*

R. Häggkvist [5] conjectured that a 2-connected bipartite graph on at most 6δ vertices is Hamiltonian. P. Ash and B. Jackson [2] showed that if

G is a 2-connected bipartite graph and each set in the bipartition has at most $3\delta - 3$ vertices, then G contains a longest cycle which is also dominating. Later, P. Ash [1, Theorem 2.1(i)] further proved that given such a graph G , $G - C$ consists entirely of components H such that $|H| \leq 2$.

As a consequence of Theorem 1.1, we obtain

COROLLARY 1.4. *A 2-connected triangle-free graph G with $|G| \leq 6\delta - 6$ contains a longest cycle which is also dominating.*

The examples in [2] show that the upper bound $6\delta - 6$ in Theorem 1.1 cannot be improved to $6\delta - 4$, and that a longest cycle in a bipartite graph on $n \geq 4\delta + 4$ vertices is not necessarily dominating.

In a forthcoming paper, Theorem 1.1 will be used in proving that every longest cycle in a 2-connected δ -regular bipartite graph on at most $6\delta - 6$ vertices is dominating.

We mention that the existence of cycles C in a graph G such that $G - C$ has only small components was studied by H. J. Veldman [8, 9], P. Fraisse [4], and J. A. Bondy and G. Fan [3], though the cycles constructed in those papers are not necessarily longest cycles, and that the existence of a (not necessarily longest) dominating cycle in a 2-connected triangle-free graph G with $|G| \leq 6\delta - 5$ can easily be deduced from a theorem of H. J. Veldman [8, Theorem 3].

2. PRELIMINARIES

Let G be a graph, u and v distinct vertices of G , and K a subgraph of G .

We shall write " $u \sim v$ " to indicate that u and v are adjacent. If H is a subgraph of G then the set $\{w \in V(K) : w \sim y \text{ for some } y \in V(H)\}$ will be denoted by $N_K(H)$ or $N_K V(H)$. Abbreviate $|N_K(v)|$ to $d_K(v)$ and $d_G(v)$ to $d(v)$. A uv -path is a path joining u and v . If $\{u, v\} \subset V(K)$ and K is connected, then a longest uv -path P with $V(P) \subset V(K)$ will be denoted by $L_K(u - v)$.

If P is a path in G and H is a component of $G - P$, then P^H denotes the longest subpath of P with terminal vertices in $N_P(H)$. For vertices a, b on P , let $P[a, b]$ denote the subpath of P with terminal vertices a and b (with the order from a to b if necessary).

Let C be a cycle in G and x, y two distinct vertices on C . After designating the orientation of C , $[x, y]$ denotes the segment (path) of C joining x and y such that the orientation of the segment from x to y is the same as that of C ; $[x, y)$ and (x, y) denote the paths $[x, y] - y$ and $[x, y] - \{x, y\}$, respectively.

LEMMA 2.1. Let $Q = Q[u, w]$ be a uw -path, in a triangle-free graph, where $u \neq w$. Let v be a vertex on Q . Then,

- (a) $|Q| \geq 2d_Q(u)$,
- (b) $|Q| \geq 2d_Q(v) - 1$,
- (c) $|Q| \geq d_Q(u) + d_Q(v)$,
- (d) $|Q| \geq 2d_Q(v)$ if $v \notin N_Q(u) \cap N_Q(w)$,
- (e) $|Q| \geq 2d_Q(v) + 1$ if $v \notin N_Q\{u, w\}$,
- (f) $|Q| \geq 2d_Q(u) + 1$ if $u \notin N_Q(w)$.

Proof. (a) can easily be checked, and (b)–(f) are consequences of (a). ■

The proof of the following lemma is partially based on that of Theorem 2.1 in [6].

LEMMA 2.2. Let a and b be distinct vertices and P a longest ab -path in the 2-connected triangle-free graph G . Each component H of $G - P$ has a vertex v such that $|P^H| \geq 2d(v) - 1$.

Proof. By induction on $|G|$. Let H be a component of $G - P$, and label $N_P(H) = \{x_1, x_2, \dots, x_s\}$ according to the order on P from a to b . Since G is 2-connected, $s \geq 2$. Since P is a longest ab -path, x_{i+1} is not the successor of x_i . Therefore, $|P^H| \geq 2s - 1 + |P[x_i, x_{i+1}]]| - 3$ for any $i < s$. Since the result is immediate if $|H| = 1$, we assume that $|H| \geq 2$.

If $G - P^H \neq H$, then the subgraph H^* of G induced by $V(H) \cup V(P^H)$ is a 2-connected triangle-free proper subgraph of G in which P^H is a longest x_1x_s -path. By the induction hypothesis, $|P^H| \geq 2d_{H^*}(v) - 1 = 2d(v) - 1$ for some $v \in V(H)$.

If H contains a cut vertex, choose a longest path $Q = Q[c_1, c_2]$ in H whose terminal vertices are cut vertices of H (may be $c_1 = c_2$). Further determine two distinct components K_1 of $H - c_1$ and K_2 of $H - c_2$ such that K_1 and K_2 contain no vertex of Q . Since G is 2-connected, $N_P(K_1) \neq \emptyset \neq N_P(K_2)$. Notice that, for $h = 1$ and 2 , the subgroup K_h^* of G induced by $V(K_h) \cup \{c_h\}$ is a triangle-free block.

Label $N_P(K_1 \cup K_2) = \{z_1, z_2, \dots, z_q\}$ according to the order on P from a to b .

First assume $q \geq 2$. There is some $j < q$ such that $z_j \in N_P(K_1)$ and $z_{j+1} \in N_P(K_2)$ or else $z_j \in N_P(K_2)$ and $z_{j+1} \in N_P(K_1)$, say $w_1 \in N_{K_1}(z_j)$ and $w_2 \in N_{K_2}(z_{j+1})$. Let Q_h be a longest $w_h c_h$ -path in H , for $h = 1, 2$. If $V(K_h) \subset V(Q_h)$ then, by Lemma 2.1(a), $|Q_h| \geq 2d_H(w_h) \geq d_H(w_h) + 1$. If $V(K_h) - V(Q_h) \neq \emptyset$ then $|Q_h| \geq 2d_{K_h^*}(v) - 1 = 2d_H(v) - 1 \geq d_H(v) + 1$ for some $v \in V(K_h^* - Q_h)$ by the induction hypothesis. In any case we can

choose a vertex $v_h \in V(K_h)$ such that $|Q_h| \geq d_H(v_h) + 1$, for $h = 1$ and 2 . Thus, $|P^H| \geq 2q - 1 + |P[z_j, z_{j+1}]| - 3 \geq 2q - 1 + |Q_1 \cup Q \cup Q_2| - 1 \geq 2q - 1 + |Q_1| + |Q_2| - 2 \geq d_P(v_1) + d_P(v_2) - 1 + d_H(v_1) + d_H(v_2) = d(v_1) + d(v_2) - 1$.

Assume, now, $q = 1$. Since $s = |N_P(H)| \geq 2$, there is some $z \in N_P(H) - \{z_1\}$. Pick $x \in N_{K_1}(z_1)$ and $y \in N_H(z)$. Then $y \notin V(K_1) \cup V(K_2)$. Let Q_1 be a longest xc_1 -path in H and Q_2 a c_1y -path in H . As we have shown, one can choose a vertex $v \in V(K_1)$ such that $|Q_1| \geq 2d_H(v) - 1$; then, $|P^H| \geq |P[z_1, z]| \geq 2 + |Q_1 \cup Q_2| \geq 2 + |Q_1| \geq 2(1 + d_H(v)) - 1 \geq 2d(v) - 1$.

Suppose, now, that $G - P^H = H$ and H has no cut vertex. Then, $a = x_1$ and $b = x_s$.

Case 1. $s \geq 3$. Since G is 2-connected and $|H| \geq 2$, there is some $x_i \in N_P(H)$ such that $|N_H\{x_i, x_{i+1}\}| \geq 2$. Let G^* be the subgraph of G induced by $V(H) \cup V(P[x_i, x_{i+1}])$. Then, G^* is a 2-connected triangle-free proper subgraph of G in which $P[x_i, x_{i+1}]$ is a longest $x_i x_{i+1}$ -path. By the induction hypothesis, $|P[x_i, x_{i+1}]| \geq 2d_{G^*}(v) - 1$ for some $v \in V(H)$. Consequently, $|P^H| \geq 2s - 1 + |P[x_i, x_{i+1}]| - 3 \geq 2(s - 2) + 2d_{G^*}(v) - 1 \geq 2dG - G^*(v) + 2d_{G^*}(v) - 1 = 2d(v) - 1$.

Case 2. $s = 2$. Since $|H| \geq 2$ and G is 2-connected, one can choose two distinct vertices $u \in N_H(a)$ and $w \in N_H(b)$ in such a way that $|L_H(u - w)|$ is maximum. Let Q be a longest uw -path in H .

First assume $|Q| = |H|$. Since $|Q| \geq 2$ and G is triangle-free, there is some $v \in V(Q)$ such that $d_P(v) \leq 1$. By Lemma 2.1(b), $|Q| \geq 2d_Q(v) - 1 = 2d_H(v) - 1$. Hence, $|P| \geq 2 + |Q| \geq 2 + 2d_H(v) - 1 \geq 2d(v) - 1$.

Assume, now, $V(H) - V(Q) \neq \emptyset$. Clearly, $|Q| \geq 3$ and hence $|P^H| \geq 5$. Form a new graph G^* by removing all the vertices of $V(P) - \{a, b\}$ from G , adding a new vertex, and joining it to a and b . Then, G^* is a 2-connected triangle-free graph and $|G^*| < |G|$. And $Q^* := aQ[u, w]b$ is a longest ab -path in G . By the induction hypothesis, $|Q^*| \geq 2d_{G^*}(v) - 1$ for some $v \in V(H - Q)$. Since $|P^H| \geq |Q^*|$ and $d_{G^*}(u) = d_G(u)$ for every $u \in V(H)$, $|P^H| \geq 2d(v) - 1$. ■

3. PROOF OF PROPOSITION 1.2

Let G be a 2-connected triangle-free graph. Suppose that for some longest cycle C' in G , there is a component H' , in $G - C'$, such that $|H'| \geq 3$. Among all pairs C', H' such that C' is a longest cycle and H' is a component of $G - C'$ with $|H'| \geq 3$, we choose one for which $|H'|$ is as small as possible, and denote it by C, H .

We shall prove this proposition by a discussion on the structure of H . First we shall discard easily the case when $|H| = 3$, then treat in two

separate proofs the case where H is 2-connected, and the case where H admits a cut-vertex.

Choose an orientation of C and fix it throughout the proof.

Since $|H| \geq 3$ and G is 2-connected, $|G| \geq 9$; assume, therefore, $\delta \geq 3$.

CLAIM 3.1. *If $|H| = 3$, then $|G| \geq 6\delta - 5$.*

Proof. Suppose $|H| = 3$. Then, H is a path (star) on three vertices, for it is triangle-free. Let v_1 and v_2 be the two terminal vertices and u the middle vertex of H . Label $N_C\{v_1, v_2\} = \{y_1, y_2, \dots, y_r\}$ in the orientation around C (the subscripts of y will be reduced modulo r). Notice that $r \geq 2$ since $\delta \geq 3$. One can easily check, using the maximality of $|C|$, that

$$|[y_i, y_{i+1}]| \geq 2 + 2d_{(y_i, y_{i+1})}(u) \quad \text{for every } y_i \quad (1)$$

and

$$|[y_i, y_{i+1}]| \geq 4 + 2d_{(y_i, y_{i+1})}(u) \quad \text{if } N_C\{y_i, y_{i+1}\} = \{v_1, v_2\}. \quad (2)$$

As G is triangle-free, $N_C(u) \cap N_C\{v_1, v_2\} = \emptyset$ and hence $d_C(u) = \sum_{i=1}^r d_{(y_i, y_{i+1})}(u)$. Accordingly, using (1) and (2),

$$\begin{aligned} |C| &= \sum_{i=1}^r |[y_i, y_{i+1}]| \\ &\geq 4 |N_C(v_1) \cap N_C(v_2)| + 2 |N_C(v_1) - N_C(v_2)| \\ &\quad + 2 |N_C(v_2) - N_C(v_1)| + 2d_C(u) \\ &= 2d_C(v_1) + 2d_C(v_2) + 2d_C(u) \\ &= 2d(v_1) + 2d(v_2) + 2d(u) - 8. \end{aligned}$$

Therefore, $|G| \geq |H| + |C| \geq 6\delta - 5$. ■

In view of Claim 3.1, we assume, from now on, that $|H| \geq 4$.

Before we check other cases, we shall develop some estimates for $|G|$ in which valences of two distinct vertices are involved.

In the following part, v_1 and v_2 will be two distinct vertices in H which are not cut vertices of H , and Q a path (containing at least two vertices) in H . In connection with v_1, v_2 , and Q , the following notation will be used if $N_C(v_1) \neq \emptyset \neq N_C(v_2)$ and $|N_C\{v_1, v_2\}| \geq 2$;

$$R_0 := \{i \mid y_i \in N_C(v_1) \cap N_C(v_2)\},$$

$$R_1 := \{i \mid y_i \in N_C(v_1) - N_C(v_2) \text{ and } y_{i+1} \in N_C(v_2)\},$$

$$R_{11} := \{i \mid \{y_i, y_{i+1}\} \subset N_C(v_1) - N_C(v_2)\},$$

and, R_2 and R_{22} are defined by interchanging v_1 and v_2 in R_1 and R_{11} . Note that these sets are disjoint and $d_C(v_h) = |R_0| + |R_h| + |R_{hh}|$ for $h = 1, 2$,

$$r_{hh} := \sum_{i \in R_{hh}} (|[y_i, y_{i+1}]| - 3) \quad \text{for } h = 1, 2,$$

$$r_h := \sum_{i \in R_h} (|[y_i, y_{i+1}]| - (|Q| + 1)) \quad \text{for } h = 0, 1, 2,$$

and

$$q := |Q| + 1 - d_H(v_1) - d_H(v_2).$$

We shall later consider several cases in which we will choose v_1, v_2 , and Q in such a way that q, r_0, r_1, r_2, r_{11} , and r_{22} are non-negative and $|H| - |Q| + 3q + r_0 + r_1 + r_2 + r_{11} + r_{22}$ is sufficiently large.

CLAIM 3.2. (a) $|[y_i, y_{i+1}]| \geq 3$ for every $i \in R_{11} \cup R_{22}$.

(b) Consequently, $r_{11}, r_{22} \geq 0$.

Proof. Let $i \in R_{hh}$ and $h = 1$ or 2 . Since C is a longest cycle, $|[y_i, y_{i+1}]| \geq 2$. If $|[y_i, y_{i+1}]| = 2$ then $C' := v_h[y_{i+1}, y_i]v_h$ is also a longest cycle. As v_h is not a cut vertex of H , $H - v_h$ is a component of $G - C'$. But $|H| > |H - v_h| = |H| - 1 \geq 3$ and hence the pair $C', H - v_h$ is better than C, H ; this contradicts the choice of C, H . Therefore, $|[y_i, y_{i+1}]| \geq 3$. ■

CLAIM 3.3. If $|Q| \leq |L_H(v_1 - v_2)|$ then

(a) $|[y_i, y_{i+1}]| \geq |Q| + 1$ for every $i \in R_0 \cup R_1 \cup R_2$, and consequently,

(b) $r_0, r_1, r_2 \geq 0$.

Proof. For each $i \in R_0 \cup R_1 \cup R_2$, we have a cycle C_i such that $V(C_i) = V([y_{i+1}, y_i]) \cup V(L_H(v_1 - v_2))$ by the definitions of R_0, R_1 , and R_2 . From $|C_i| \leq |C|$ follows $|[y_i, y_{i+1}]| \geq |L_H(v_1 - v_2)| + 1$. ■

CLAIM 3.4. (a) If

(i) $|C| \geq 2|Q| + 2$,

(ii) $|Q| \geq 2d_H(v_h) - 1$ for $h = 1, 2$, and

(iii) $N_C(v_h) = \emptyset$, then

$$|C| + |H| \geq 6d(v_h) - 1.$$

(b) If $|N_C\{v_1, v_2\}| \leq 1$, then

$$|C| + |H| \geq 3d(v_1) + 3d(v_2) - 7 + 3q + (|H| - |Q|) + (|C| - 2|Q| - 2).$$

(c) If

(i) $N_C(v_1) \neq \emptyset \neq N_C(v_2)$, and

(ii) $v_1 \sim v_2$, then

$$|C| + |H| \geq 3d(v_1) + 3d(v_2) - 7 + 3q + (|H| - |Q|) + (r_1 + r_2 + r_{11} + r_{22}).$$

(d) If

(i) $N_C(v_1) \neq \emptyset \neq N_C(v_2)$.

(ii) $|N_C\{v_1, v_2\}| \geq 2$, and

(iii) $|Q| \geq 5$,

then

$$\begin{aligned} |C| + |H| &\geq 3d(v_1) + 3d(v_2) - 13 + 3q + (|H| - |Q|) \\ &\quad + (r_0 + r_1 + r_2 + r_{11} + r_{22}). \end{aligned}$$

Proof. (a) Since $|H| \geq |Q|$ then $|C| + |H| \geq 3|Q| + 2$ by (i). By (iii), $d_H(v_h) = d(v_h)$. Then, the conclusion follows by (ii).

(b) Trivially, $|C| + |H| \equiv 3|Q| + 2 + (|H| - |Q|) + (|C| - 2|Q| - 2)$. By the definition of q , $3|Q| + 2 = 3d_H(v_1) + 3d_H(v_2) - 1 + 3q$. For $h = 1$ and 2 , $d_H(v_h) \geq d(v_h) - 1$ since $|N_C(v_h)| \leq 1$. Hence, the assertion.

(c) By (i) and (ii), $R_0 = \emptyset$ (since G is triangle-free), $|N_C\{v_1, v_2\}| \geq 2$, and $|R_1| + |R_2| \geq 2$. Since $|R_0| = 0$ then $d_C(v_h) = |R_h| + |R_{hh}|$ for $h = 1, 2$. By the definitions of r_1, r_2, r_{11} , and r_{22} , we have $|C| = \sum_{i=1}^r |y_i, y_{i+1}| = (|Q| + 1)(|R_1| + |R_2|) + 3(|R_{11}| + |R_{22}|) + (r_1 + r_2 + r_{11} + r_{22})$. Consequently,

$$\begin{aligned} |C| + |H| &= 3|Q| + 2 + (|Q| + 1)(|R_1| + |R_2| - 2) + 3(|R_{11}| + |R_{22}|) \\ &\quad + (|H| - |Q|) + (r_1 + r_2 + r_{11} + r_{22}). \end{aligned}$$

By the definition of q , we have $3|Q| + 2 = 3d_H(v_1) + 3d_H(v_2) - 1 + 3q$. From $|Q| + 1 \geq 3$ and $|R_1| + |R_2| \geq 2$, it follows that $(|Q| + 1)(|R_1| + |R_2| - 2) \geq 3(|R_1| + |R_2| - 2)$. Further, $3(|R_1| + |R_2| - 2) + 3(|R_{11}| + |R_{22}|) = 3(|R_1| + |R_{11}|) + 3(|R_2| + |R_{22}|) - 6 = 3d_C(v_1) + 3d_C(v_2) - 6$. We then obtain the required inequality.

(d) From (i) and (ii), it follows that $|R_0| + |R_1| + |R_2| \geq 2$. By the definitions of $r_0, r_1, r_2, r_{11}, r_{22}$, we have

$$\begin{aligned} |C| &= (|Q| + 1)(|R_0| + |R_1| + |R_2|) + 3(|R_{11}| + |R_{22}|) \\ &\quad + (r_0 + r_1 + r_2 + r_{11} + r_{22}) \end{aligned}$$

and then

$$\begin{aligned} |C| + |H| &= 3|Q| + 2 + (|Q| + 1)(|R_0| + |R_1| + |R_2| - 2) + 3(|R_{11}| + |R_{22}|) \\ &\quad + (|H| - |Q|) + (r_0 + r_1 + r_2 + r_{11} + r_{22}). \end{aligned}$$

By the definition of q , $3|Q| + 2 = 3d_H(v_1) + 3d_H(v_2) - 1 + 3q$. From $|Q| + 1 \geq 6$ (by (iii)) and $|R_0| + |R_1| + |R_2| \geq 2$, it follows that $(|Q| + 1)(|R_0| + |R_1| + |R_2| - 2) \geq 6(|R_0| + |R_1| + |R_2| - 2)$. Recall also that $|R_0| + |R_h| + |R_{hh}| = d_C(v_h)$ for $h = 1, 2$. We now reach the inequality. ■

Our next task is to find v_1, v_2 , and Q which produce a sufficiently large $(|H| - |Q|) + 3q + (r_0 + r_1 + r_2 + r_{11} + r_{22})$. The method of choosing v_1, v_2 , and Q is based on the method that H. A. Jung used to prove Lemma 3.2 in [6].

I. Suppose that H Is 2-Connected

Label $N_C(H) = \{x_1, x_2, \dots, x_s\}$ in the orientation around C (the subscripts will be reduced modulo s).

Let $I := \{i: |N_H\{x_i, x_{i+1}\}| \geq 2\}$. Since $|H| > 1$ and G is 2-connected, then $|I| \geq 2$. One can choose, for any $i \in I$, a longest path $Q_i = Q_i[u_i, w_i]$ in H such that $u_i \in N_H(x_i)$ and $w_i \in N_H(x_{i+1})$. Since C is a longest cycle, $|[x_i, x_{i+1})| \geq |Q_i| + 1$ for each $i \in I$. Pick $j \in I$ such that $|Q_j| \leq |Q_i|$ for every $i \in I$. Note that $|Q_j| \geq 3$ since $|H| > 2$ and H is 2-connected.

CLAIM 3.5. (a) If $x_k \neq x_l$ and $|N_H\{x_k, x_l\}| \geq 2$ then $|[x_k, x_l)| \geq |Q_j| + 1$.

(b) In particular, $|C| \geq 2|Q_j| + 2$, and, if $Q = Q_j$, then

(c) $|[y_i, y_{i+1})| \geq |Q| + 1$ for every $i \in R_0 \cup R_1 \cup R_2$, and consequently,

(d) $r_0, r_1, r_2 \geq 0$.

Proof. Suppose $x_k \neq x_l$ and $|N_H\{x_k, x_l\}| \geq 2$. Then, there is some $x_m \in V([x_k, x_l))$ such that $|N_H\{x_m, x_{m+1}\}| \geq 2$. By the definition of Q_j , we have $|Q_m| \geq |Q_j|$. Hence, $|[x_k, x_l)| \geq |[x_m, x_{m+1})| \geq |Q_m| \geq |Q_j|$.

Since $|N_H\{x_j, x_{j+1}\}| \geq 2$, we obtain $|[x_j, x_{j+1})|, |[x_{j+1}, x_j)| \geq |Q_j| + 1$. Hence, $|C| \geq 2|Q_j| + 2$.

By the definitions of R_0, R_1 , and R_2 , (c) is a direct consequence of (a),

and, by the definitions of r_0, r_1 , and r_2 , (d) is a direct consequence of (c). ■

Case 1. $|Q_j| = 3$. Since Q_j is a longest $u_j w_j$ -path in H and H is 2-connected, $N_H(v) = \{u_j, w_j\}$ for every $v \in V(H) - \{u_j, w_j\}$.

If $|H| - |Q_j| = 1$, then H is a cycle on four vertices. Choose any two adjacent vertices of H as v_1 and v_2 , and $L_H(v_1 - v_2)$ as Q . Since $\delta > 2 = d_H(v_1) = d_H(v_2)$ then $N_C(v_1) \neq \emptyset \neq N_C(v_2)$. And $q = 5 - d_H(v_1) - d_H(v_2) = 1$. Accordingly, $|G| \geq 6\delta - 4$ by Claims 3.4(c), 3.2(b), and 3.3(b).

If $|H| - |Q_j| \geq 2$, choose any two distinct vertices from $V(H) - \{u_j, w_j\}$ as v_1 and v_2 , and $L_H(v_1 - v_2)$ as Q . Then $|Q| = 5$. By Claim 3.5(b), $|C| \geq 2|Q_j| + 2 = 8$ and then $|G| \geq 13$. Assume, therefore, $\delta \geq 4$. Then $d_C(v_1), d_C(v_2) \geq 2$, and $|G| \geq 6\delta - 13 + (3q + r_0 + r_1 + r_2 + r_{11} + r_{22})$ by Claim 3.4(d). As $\{u_j, w_j\} = N_H(v_1) \cap N_H(v_2)$ and G is triangle-free, $\{x_j, x_{j+1}\} \cap N_C\{v_1, v_2\} = \emptyset$. Suppose $\{x_j, x_{j+1}\} \subset V((y_i, y_{i+1}))$. Since $|L_H(u_j - v_h)| = |L_H(w_j - v_h)| = 4$ for $h = 1, 2$ and C is a longest cycle, $|[y_i, y_{i+1}]| = |[y_i, x_j]| + |[x_j, x_{j+1}]| + |[x_{j+1}, y_{i+1}]| \geq 14 = (|Q| + 1) + 8 > 3 + 8$; hence, by Claims 3.2(a), 3.3(a), and by the definitions of r_0, r_1, r_2, r_{11} , and r_{22} , we obtain $r_0 + r_1 + r_2 + r_{11} + r_{22} \geq 8$. And $q = 2$ since $|Q_j| = 5$ and $d_H(v_1) = d_H(v_2) = 2$. Hence, $|G| \geq 6\delta + 1$.

Case 2. $|Q_j| = 4$. Label $Q_j = u_j u w w_j$. By Claim 3.5(b), $|C| \geq 2|Q_j| + 2 \geq 10$ which implies that $|G| \geq 14$. We, therefore, assume $\delta \geq 4$.

Suppose $u \in N_{Q_j}(K_1)$ and $w \in N_{Q_j}(K_2)$ where K_1 and K_2 are components of $H - Q_j$. Since Q_j is a longest $u_j w_j$ -path in H and H is 2-connected, $K_1 \neq K_2$, $N_{Q_j}(K_1) = \{u, w_j\}$, and $N_{Q_j}(K_2) = \{u_j, w\}$. But, we then obtain a $u_j w_j$ -path which passes through K_2, w, u , and K_1 , contradicting the fact that Q_j is a longest $u_j w_j$ -path in H . We therefore have $N_{H-Q_j}(u) = \emptyset$ or $N_{H-Q_j}(w) = \emptyset$. Without loss of generality let $N_{H-Q_j}(u) = \emptyset$. Then, $d_H(u) = 2$ because H is triangle-free. Choose u as v_1 .

If $N_{H-Q_j}(w) = \emptyset$, choose w as v_2 and Q_j as Q . In this subcase, $q = 1$ and hence Claims 3.4(c), 3.5(d), and 3.2(b) apply to yield $|G| \geq 6\delta - 4$.

Let $w \in N_{Q_j}(K)$ where K is a component of $H - Q_j$. Since H is 2-connected and Q_j is a longest $u_j w_j$ -path in H , $N_{Q_j}(K) = \{u_j, w\}$ and $|K| = 1$. Choose the vertex of K as v_2 , and the path $v_2 u_j v_1 w w_j$ as Q . As $d_C(v_1), d_C(v_2) \geq 2$ (since $\delta \geq 4$ and $d_H(v_1) = d_H(v_2) = 2$) and $|Q_j| = 5$, $|G| \geq 6\delta - 13 + (3q + r_0 + r_1 + r_2 + r_{11} + r_{22})$ by Claim 3.4(d). Since $u_j \in N_H(v_1) \cap N_H(v_2)$ and $u_j \sim x_j, x_j \notin N_C\{v_1, v_2\}$. Suppose $x_j \in V((y_i, y_{i+1}))$. Since H is 2-connected and triangle-free and since Q_j is a longest $u_j w_j$ -path in H , there is a $u_j w_j$ -path in $H - \{v_2, v_1, w\}$. Hence, $|L_H(v_1 - v_2)| \geq 5 = |Q_j|$. Moreover, by Claim 3.5(a), $|[y_i, y_{i+1}]| \geq |[y_i, x_j]| + |[x_j, x_{j+1}]| \geq 2|Q_j| + 2 = 10 = (|Q_j| + 1) + 4 > 3 + 4$; hence, by Claims 3.2(a) and 3.3(a), $r_0 + r_1 + r_2 + r_{11} + r_{22} \geq 4$. And, $q = 2$ since $|Q_j| = 5$ and $d_H(v_1) = d_H(v_2) = 2$. Therefore, $|G| \geq 6\delta - 3$.

Case 3. $|Q_j| \geq 5$ and $|H| = |Q_j|$. Choose u_j as v_1 , the successor of u_j on $Q_j[u_j, w_j]$ as v_2 , and Q_j as Q . Since $|Q_j| \geq 2d_H(v_2) - 1$ by Lemma 2.1(b), if $N_C(v_2) = \emptyset$ then $|G| \geq 6d(v_2) - 1$ by Claims 3.4(a) and 3.5(b). If $N_C(v_2) \neq \emptyset$ then we obtain $|G| \geq 6\delta - 4$ by using Claims 3.4(c), 3.5(d), and 3.2(b) because $q \geq 1$ (by Lemma 2.1(c)), $N_C(v_1) \neq \emptyset$, and $v_1 \sim v_2$.

Case 4. $|Q_j| \geq 5$ and $H - Q_j$ has distinct components K_1 and K_2 . Choose, for $h = 1$ and 2 , $v_h \in V(K_h)$ such that $|Q_j^{K_h}| \geq 2d_H(v_h) - 1$ by Lemma 2.2. Choose Q_j as Q . Then, $|H| - |Q| \geq 2$ and $q = \frac{1}{2}(2|Q| + 2 - 2d_H(v_1) - 2d_H(v_2)) \geq 0$.

In view of Claims 3.4(a), 3.4(b), and 3.5(b) we assume that $N_C(v_1) \neq \emptyset \neq N_C(v_2)$ and $|N_C\{v_1, v_2\}| \geq 2$. Then, by Claim 3.4(d), $|G| \geq 6\delta - 11 + (3q + r_0 + r_1 + r_2 + r_{11} + r_{22})$. As $r_0, r_1, r_2, r_{11}, r_{22} \geq 0$ (by Claims 3.5(d) and 3.2(b)) and $q \geq 0$, it suffices to show that $q \geq 2$, or, $q \geq 1$ and $r_{11} \geq 3$, or $r_0 + r_1 - r_2 + r_{11} + r_{22} \geq 6$.

Case 4.1. $x_j \in N_C(v_1)$ and $x_{j+1} \in N_C(v_2)$ (or $x_j \in N_C(v_2)$ and $x_{j+1} \in N_C(v_1)$). By the choice of Q_j , $u_j \notin N_{Q_j}(K_1)$ and $w_j \notin N_{Q_j}(K_2)$. Moreover, if $w_j \in N_{Q_j}(K_1)$, then u_j and the successor of u_j on $Q_j[u_j, w_j]$ cannot be contained in $N_{Q_j}(K_2)$; then, $|Q_j| \geq 3 + |Q_j^{K_2}| \geq 2d_H(v_2) + 2$ and $|Q_j| \geq 1 + |Q_j^{K_1}| \geq 2d_H(v_1)$, and hence $q \geq 2$. Similarly, if $u_j \in N_{Q_j}(K_2)$ then $q \geq 2$. If $\{u_j, w_j\} \cap N_{Q_j}(K_1 \cup K_2) = \emptyset$ then $|Q_j| \geq 2 + |Q_j^{K_h}| \geq 2d_H(v_h) + 1$ for $h = 1$ and 2 ; thus, $q \geq 2$ again.

Case 4.2. $\{x_j, x_{j+1}\} \subset N_C(v_1) - N_C(v_2)$ (or $\subset N_C(v_2) - N_C(v_1)$). Then, $x_j = y_i$ and $x_{j+1} = y_{i+1}$ for some $i \in R_{11}$. By Claim 3.5(a), $|[x_j, x_{j+1}]| \geq |Q_j| + 1 \geq 6$. Accordingly, $|[y_i, y_{i+1}]| - 3 \geq 3$. This and Claim 3.2(a) together imply that $r_{11} \geq 3$. Moreover, by the choice of Q_j , $\{u_j, w_j\} \cap N_{Q_j}(K_1) = \emptyset$ and hence $|Q_j| \geq |Q_j^{K_1}| + 2 \geq 2d_H(v_1) + 1$; thus $q \geq 1$.

Case 4.3. $x_{j+1} \notin N_C\{v_1, v_2\}$ (or $x_j \notin N_C\{v_1, v_2\}$). Suppose $\{x_j, x_{j+1}\} \subset V([y_i, y_{i+1}])$. By Claim 3.5(a), $|[y_i, y_{i+1}]| \geq |[x_j, x_{j+1}]| + |[x_{j+1}, y_i]| \geq 2|Q_j| + 2 \geq (|Q_j| + 1) + 6 > 3 + 6$; this and Claims 3.2(a), 3.5(c) together imply that $r_0 + r_1 + r_2 + r_{11} + r_{22} \geq 6$.

Case 5. $|Q_j| \geq 5$ and $H - Q_j$ has only one component K . Choose Q_j as Q . Then, $|H| - |Q| \geq 1$.

Case 5.1. $u_j \in N_{Q_j}(K)$ (or $w_j \in N_{Q_j}(K)$). Choose some vertex $v_1 \in V(K)$ such that $|Q_j^K| \geq 2d_H(v_1) - 1$ by Lemma 2.2. Choose the successor of u_j on $Q_j[u_j, w_j]$ as v_2 . By Lemma 2.1(b), $|Q_j| \geq 2d_H(v_2) - 1$. Then, $q \geq 0$.

In view of Claims 3.4(a) and 3.5(b), we assume that $N_C(v_h) \neq \emptyset$ for $h = 1$ and 2 . Note that $x_j \notin N_C(v_1)$ by the choice of Q_j , and that $x_j \notin N_C(v_2)$ since $v_2 \sim u_j$ and $u_j \sim x_j$.

First assume $|N_C\{v_1, v_2\}| = 1$; that is, $N_C\{v_1, v_2\} = \{y_1\}$. Then, by Claim 3.4(b), $|G| \geq 6\delta - 6 + (3q + |C| - 2|Q| - 2)$. Suppose $y_1 = x_{j+1}$.

Then, $w_j \notin N_C(K)$ due to the choice of Q_j ; hence $|Q_j| \geq |Q_j^K| + 1 \geq 2d_H(v_1)$ which yields $q \geq 1$. Since $|C| - 2|Q| - 2 \geq 0$ (by Claim 3.5(b)) and $q \geq 1$, $|G| \geq 6\delta - 3$. Suppose, $y_1 \neq x_{j+1}$. By Claim 3.5(a), $|C| = |[x_{j+1}, Y_i]| + |[y_i, x_j]| + |[x_j, x_{j+1}]| \geq 3|Q_j| + 3 \geq (2|Q| + 2) + 6$. As $|C| - 2|Q| - 2 \geq 6$ and $q \geq 0$, $|G| \geq 6\delta$.

Assume, now, $N_C(v_1) \neq \emptyset \neq N_C(v_2)$ and $|N_C\{v_1, v_2\}| \geq 2$. Suppose $x_j \in V((y_i, y_{i+1}))$ (it was shown that $x_j \notin N_C\{v_1, v_2\}$). By Claim 3.5(b),

$$\begin{aligned} |[y_i, x_{j+1}]| &\geq |[y_i, x_j]| + |[x_j, x_{j+1}]| \geq 2|Q_j| + 2 \\ &\geq (|Q| + 1) + 6 > 3 + 6. \end{aligned} \quad (3)$$

If $x_{j+1} \in N_C(v_1)$, then $w_j \notin N_{Q_j}(K)$ by the choice of Q_j ; hence $|Q_j| \geq |Q_j^K| + 1 \geq 2d_H(v_1)$ and $q \geq 1$. If $x_{j+1} \in N_C(v_2)$ then $v_2 \notin N_H(w_j)$ (since G is triangle-free) and hence $|Q| \geq 2d_H(v_2)$ (by Lemma 2.1(d)) from which $q \geq 1$ follows. As $q \geq 1$, $|H| - |Q| \geq 1$, and $r_0 + r_1 + r_2 + r_{11} + r_{22} \geq 6$ (by (3) and by Claims 3.2(a), 3.5(c)), $|G| \geq 6\delta - 3$ by Claim 3.4(d).

Suppose $x_{j+1} \notin N_C\{v_1, v_2\}$. Then $|[x_{j+1}, y_{i+1}]| \geq |Q_j| + 1 \geq 6$ by Claim 3.5(a). This and (3) together imply that $|[y_i, y_{i+1}]| \geq (|Q_j| + 1) + 12 > 3 + 12$; that is, $r_0 + r_1 + r_2 + r_{11} + r_{22} \geq 12$. By Claim 3.4(d), $|G| \geq 6\delta$.

Case 5.2. $\{u_j, w_j\} \cap N_{Q_j}(K) = \emptyset$. Choose u_j as v_1 and w_j as v_2 . By Lemma 2.1(a), $|Q| \geq 2d_H(v_h)$ for $h = 1$ and 2 ; hence, $q \geq 1$. If $v_1 \sim v_2$ then $|G| \geq 6\delta - 3$ by Claims 3.4(c), 3.2(b), and 3.5(d).

Assume $v_1 \notin N_H(v_2)$. Then, by Lemma 2.1(f), $|Q| \geq 2d_H(v_h) + 1$ for $h = 1, 2$; hence, $q \geq 2$. Since $x_j \sim v_1$ and $x_{j+1} \sim v_2$, $x_j = y_i$ and $x_{j+1} = y_{i+1}$ for some $i \in R_0 \cup R_1$. By Claim 3.5(c), $|[y_i, y_{i+1}]| \geq |Q| + 1$. If $|[y_i, y_{i+1}]| > |Q| + 1$ then $r_0 + r_1 \geq 1$ and $|G| \geq 6\delta - 5$ by Claims 3.4(d), 3.5(d), and 3.2(b).

Suppose $|[y_i, y_{i+1}]| = |Q| + 1$. Then, $[y_{i+1}, y_i]$ and Q form a longest cycle C' in G and the path (y_i, y_{i+1}) is contained in a component H' of $G - C'$. According to the choice of C, H , we have $|H'| \geq |H| \geq |Q| + 1$. But $|H'| = |Q|$. Hence $|H' - (y_i, y_{i+1})| \geq 1$. As $V(H' - (y_i, y_{i+1})) \subset V(G - (C \cup H))$, we have $|G - (C \cup H)| \geq 1$ and then $|G| \geq 6\delta - 5$ by Claims 3.4(d), 3.5(d), and 3.2(b).

II. Suppose that H Has a Cut Vertex

Choose two cut vertices c_1 and c_2 of H in such a way that the distance between them is maximum (maybe $c_1 = c_2$). For $h = 1$ and 2 , choose a component K_h of $H - c_h$ which does not contain a cut vertex of H . Let the choice be taken so that $K_1 \neq K_2$ and $|K_1| + |K_2|$ is maximum. Denote by K_h^* the subgraph of H induced by $V(K_h) \cup \{c_h\}$, for $h = 1$ and 2 . Notice that K_h^* is a triangle-free block. Label $N_C(K_1 \cup K_2) = \{t_1, t_2, \dots, t_k\}$ in the

orientation around C (the subscripts will be reduced modulo k). Since G is 2-connected, $N_C(K_1) \neq \emptyset \neq N_C(v_2)$ and hence $k \geq 1$.

Suppose, first, that $k = 1$. Without loss of generality let $|K_1| \leq |K_2|$. Choose a vertex $w_1 \in N_{K_1}(t_1)$ and a longest $w_1 c_1$ -path Q_1 in H . Since G is 2-connected there is some $x \in N_C(H) - \{t_1\}$. Suppose $y \in N_H(x)$. Then $y \notin V(K_1 \cup K_2)$. Let Q' be a $c_1 y$ -path in H . Clearly $Q_1 \cup Q'$ form a $w_1 y$ -path in H . Since C is a longest cycle in G , $|[x, t_1]| \geq 1 + |Q_1 \cup Q'| \geq 1 + |Q_1|$, and $|[t_1, x]| \geq 1 + |Q_1|$ too. Hence, $|C| + |H| \geq 2 + 2|Q_1| + |K_1^* \cup K_2^*| \geq 4|Q_1| + 1$. If $|Q_1| = |K_1^*|$, then $|Q_1| \geq 2d_{K_1^*}(w_1) = 2d_H(w_1)$ by Lemma 2.1(a); if $|Q_1| \neq |K_1^*|$ we can choose a vertex $v \in V(K_1^* - Q_1)$ such that $|Q_1| \geq 2d_{K_1^*}(v) - 1 = 2d_H(v) - 1$ by Lemma 2.2. Hence, $|G| \geq 4|Q_1| + 1 \geq 8d_H(v) - 3 \geq 8d(v) - 11 \geq 6d(v) - 5$ for some $v \in V(K_1)$.

Suppose now that $k \geq 2$. Let $I := \{i: t_i \in N_C(K_1) \text{ and } t_{i+1} \in N_C(K_2), \text{ or } t_i \in N_C(K_2) \text{ and } t_{i+1} \in N_C(K_1)\}$. Since $k \geq 2$, $|I| \geq 2$. For each $i \in I$, one can determine a longest $u_i w_i$ -path in H such that $u_i \in N_{K_1}(t_i)$ and $w_i \in N_{K_2}(t_{i+1})$, or, $u_i \in N_{K_1}(t_{i+1})$ and $w_i \in N_{K_2}(t_i)$. Pick some $j \in I$ such that $|Q_j| \leq |Q_i|$ for every $i \in I$. Assume, without loss of generality, that $u_j \in N_{K_1}(t_j)$ and $w_j \in N_{K_2}(t_{j+1})$. Clearly, Q_j passes through c_1 and c_2 , and $Q_j[u_j, c_1]$, $Q_j[c_1, c_2]$, $Q_j[c_2, w_j]$ are longest paths (in H) joining mentioned terminal vertices. Denote these subpaths of Q_j by Q^1 , Q^0 , and Q^2 , respectively.

The following claim is analogous to Claim 3.5(b)–(d) and we therefore omit the proof.

CLAIM 3.6. (a) $|C| \geq 2|Q_j| + 2$, and, if $Q = Q_j$ and $v_h \in V(K_h)$ for $h = 1$ and 2, then

(b) $|[y_i, y_{i+1}]| \geq |Q| + 1$ for every $i \in R_0 \cup R_1 \cup R_2$, and consequently

(c) $r_0, r_1, r_2 \geq 0$. ■

Case 1. $|Q_j| \geq 5$. Choose Q_j as Q . We shall obtain $|G| \geq 6\delta - 5$ by Claims 3.4(a), (b), (d), 3.6(a), 3.2(b), and 3.6(c) if we can find vertices $v_1 \in V(K_1)$ and $v_2 \in V(K_2)$ such that (i) $|Q| \geq 2d_H(v_h) - 1$ for $h = 1$ or 2, and (ii) $q \geq 3$, or, $q \geq 2$ and $|H| - |Q| \geq 2$.

If $|Q^1| = |K_1^*|$ choose u_j as v_1 and if $|Q^1| \neq |K_1^*|$ choose a vertex $v_1 \in V(K_1 - Q^1)$ such that $|Q^1| \geq 2d_H(v_1) - 1$ by Lemma 2.2. If $|Q^2| = |K_2^*|$ choose w_j as v_2 and if $|Q^2| \neq |K_2^*|$ choose a vertex $v_2 \in V(K_2 - Q^2)$ such that $|Q^2| \geq 2d_H(v_2) - 1$ by Lemma 2.2. Then, $|Q^h| \geq 2d_H(v_h)$ by Lemma 2.1(a) in case $|Q^h| = |K_h^*|$, for $h = 1, 2$.

If $d_H(v_1) = 1$ then $|Q| \geq 2d_H(v_1) + 3$ and $|Q| \geq |Q^2| + 1 \geq 2d_H(v_2)$; hence $q \geq 3$. Similarly, $q \geq 3$ if $d_H(v_2) = 1$.

Suppose $d_H(v_1), d_H(v_2) \geq 2$. Then, $|Q^h| \geq 3$ for $h = 1$ and 2. If $v_1 = u_j$, then $|Q| \geq |Q^1| + 2 \geq 2d_H(v_1) + 2$ by Lemma 2.1(a) while $|Q| \geq |Q^2| + 2 \geq 2d_H(v_2) + 1$; hence $q \geq 3$. Similarly, $q \geq 3$ if $v_2 = w_j$. If $v_1 \neq u_j$ and $v_2 \neq w_j$

then $|H| - |Q| \geq 2$, and $|Q| \geq 2 + |Q^h| \geq 2d_H(v_h) + 1$ (for $h = 1, 2$) which yields $q \geq 2$.

Case 2. $|Q_j| = 4$, $|Q^0| = 1$. Then, $|Q^1| = 3$ and $|Q^2| = 2$, or $|Q^1| = 2$ and $|Q^2| = 3$. Assume, without loss of generality, $|Q^1| = 3$ and $|Q^2| = 2$. Since K_1^* is a triangle-free block and Q_1 is a longest $u_j c_1$ -path, $V(K_1 - Q^1) \neq \emptyset$ and $N_H(v) = \{u_j, c_1\}$ for every $v \in V(K_1) - \{u_j, c_1\}$. Choose a vertex of $V(K_1) - \{u_j, c_1\}$ as v_1 and w_j as v_2 and $L_H(v_1 - v_2)$ as Q . Then, $|Q| = 5 = d_C(v_1) + d_C(v_2) + 2$ and hence $q = 3$. Since $d_C(v_1) \geq 1$, $d_C(v_2) \geq 2$ (as $\delta \geq 3$), $|Q| = 5$, and $q = 3$, then Claims 3.4(d), 3.6(c), and 3.2(b) apply to yield $|G| \geq 6\delta - 4$.

Case 3. $|Q_j| = 4$, $|Q^0| \neq 1$. By the choice of c_1, c_2, K_1, K_2 , and Q_j , then $c_1 \sim c_2$, and, for every $v \in V(H) - \{c_1, c_2\}$, $N_H(v)$ is either $\{c_1\}$ or $\{c_2\}$. Without loss of generality let $d_H(c_1) \leq d_H(c_2)$. Then, $|H| \geq 2d_H(c_1)$.

Let x, y be two distinct vertices of $N_C\{u_j, c_1, w_j\}$. By the maximality of $|C|$, we find that $|[x, y]| \geq 2$ and that $|[x, y]| \geq 4$ in case $x \in (N_C(u_j) \cap N_C(w_j)) \cup (N_C(w_j) \cap N_C(c_1))$. And, $N_C(u_j) \cap N_C(c_1) = \emptyset$ as $u_j \sim c_1$. Using this observation, we obtain $|C| \geq 2d_C(u_j) + 2d_C(w_j) + 2d_C(c_1) \geq 4\delta + 2d_C(c_1) - 4$; as $|H| \geq 2d_H(c_1)$ we further obtain $|G| \geq 6\delta - 4$.

Case 4. $|Q_j| = 3$ (i.e., $c_1 = c_2$, $V(K_1) = \{u_j\}$, and $V(K_2) = \{w_j\}$). By the choice of c_1 and c_2 , vertex c_1 is the only cut vertex of H . By the maximality condition in the choice of K_1 and K_2 , therefore H is a star.

Since C is a longest cycle,

$$|[t_i, t_{i+1}]| \geq 4 + 2d_{(t_i, t_{i+1})}(c_1) \geq 5 + d_{(t_i, t_{i+1})}(c_1)$$

for every i such that $N_{(t_i, t_{i+1})}(c_1) \neq \emptyset$, and, $|[t_i, t_{i+1}]| \geq 4$ if $\{u_j, w_j\} \subset N_H\{t_i, t_{i+1}\}$. For the remaining segments $[t_i, t_{i+1})$, Claim 3.2(a) (with $\{u_j, w_j\} = \{v_1, v_2\}$) yields $|[t_i, t_{i+1}]| \geq 3$. Divide the set $\{t_i \in N_C(u_j) \cap N_C(w_j) : N_{(t_i, t_{i+1})}(c_1) = \emptyset\}$ into two sets $A := \{t_i \in N_C(u_j) \cap N_C(w_j) : |[t_i, t_{i+1}]| = 4\}$ and $B := \{t_i \in N_C(u_j) \cap N_C(w_j) : |[t_i, t_{i+1}]| \geq 5\}$, and put $D := \{t_i \in N_C(u_j) \cap N_C(w_j) : N_{(t_i, t_{i+1})}(c_1) \neq \emptyset\}$. Then,

$$\begin{aligned} |C| \geq & 4|A| + 5|B| + 5|D| + 3|N_C(u_j) - N_C(w_j)| \\ & + 3|N_C(w_j) - N_C(u_j)| + d_C(c_1). \end{aligned} \quad (4)$$

We next show that

$$|G - (C \cup H)| \geq |A|. \quad (5)$$

As in Case 5.2 of Part I, one can see that, for each $t_i \in A$, $N_{G-(C \cup H)}((t_i, t_{i+1})) \neq \emptyset$. Assume that x on (t_i, t_{i+1}) and y on (t_l, t_{l+1}) have a common neighbour $v \in V(G - (C \cup H))$, where t_i and t_l are distinct

elements of A . Then, the paths $[y, t_i]$, $t_i u_j c_1 w_j t_l$, $[x, t_l]$, and xvy form a cycle C' . From $|C'| \leq |C|$ follows $|[t_i, x]| + |[t_l, y]| \geq 4$. Similarly, $|[x, t_{i+1}]| + |[y, t_{l+1}]| \geq 2$. We then obtain $|[t_i, t_{i+1}]| + |[t_l, t_{l+1}]| \geq 8$, a contradiction. Hence (5).

By (4) and (5), $|G - H| \geq 5(|A| + |B| + |D|) + 3|N_C(u_j) - N_C(w_j)| + 3|N_C(w_j) - N_C(u_j)| + d_C(c_1)$. From $|A| + |B| + |D| = |N_C(u_j) \cap N_C(w_j)|$ follows $|G - H| \geq 3d_C(u_j) + 2d_C(w_j) + d_C(c_1)$. Since $d_H(u_j) = d_H(w_j) = 1$ and $|H| = d_H(c_1) + 1$, $|G| \geq 3d(u_j) + 2d(w_j) + d(c_1) - 4 \geq 6\delta - 4$.

4. PROOF OF PROPOSITION 1.3

The observation of H. A. Jung used to prove Lemma 2.7 in [7] leads to this proof.

Let G be a triangle-free graph and C a longest cycle in it. Suppose that $G - C$ contains two distinct components H and H^* with $|H| = |H^*| = 2$. Since the result is trivial for $\delta = 1$, we assume $\delta \geq 2$.

Choose an orientation of C and fix it. Label $N_C(H) = \{x_1, x_2, \dots, x_s\}$ in the orientation around C . The subscripts will be reduced modulo s . Define

$$X := \{i: V([x_{i-1}, x_i]) \cap N_C(H^*) \neq \emptyset\}$$

and

$$Y := \{i: V([x_{i-1}, x_i]) \cap N_C(H^*) = \emptyset\}.$$

For each $i \in X$, denote by y_i the last vertex of $N_C(H^*)$ on the path $[x_{i-1}, x_i]$. Pick $j \in X$ such that $|[y_j, x_j]| \leq |[y_i, x_i]|$ for every $i \in X$. Define, for every i ,

$$\sigma_i := |[x_{i-1}, x_i]| - 2|N_C(H^*) \cap V([x_{i-1}, x_i])|.$$

As C is a longest cycle, any two consecutive vertices cannot both be in $N_C(H)$ or $N_C(H^*)$. Consequently, $\sigma_i \geq 2$ for every $i \in Y$ because $\sigma_i = |[x_{i-1}, x_i]|$ by the definition of Y , and $\sigma_i \geq |[y_i, x_i]| - 1$ for every $i \in X$.

We let $V(H) = \{a, b\}$, $V(H^*) = \{u, v\}$, $N_H\{x_j\} = \{a\}$, and $N_{H^*}\{y_j\} = \{u\}$.

Case 1. $|[y_j, x_j]| \geq 2$. Then, $\sigma_i \geq 1$ for every $i \in X$. Hence, $|C| = \sum_{i=1}^s |[x_i, x_{i+1}]| \geq 2|N_C(H^*)| + |X| + 2|Y| \geq 2|N_C(H^*)| + |N_C(H)| \geq 6\delta - 6$ and $|G| \geq 6\delta - 2$.

Case 2. $|[y_j, x_j]| = 1$. Then $\sigma_i \geq 0$ for every $i \in X$. Take any $i \in X - \{j\}$. Let L_i be the longest $x_j x_i$ -path and M_i the longest $y_j y_i$ -path such that $V(L_i) - \{x_j, x_i\} \subset V(H)$ and $V(M_i) - \{y_j, y_i\} \subset V(H^*)$. Then, $[x_i, y_j]$, M_i ,

$[x_j, y_i]$, and L_i form a cycle C' . From $|C'| \leq |C|$ follows $|(y_i, x_i)| \geq |L_i| + |M_i| - 5$; hence $\sigma_i \geq |L_i| + |M_i| - 6$. Accordingly,

$$\sigma_i \geq 1 \quad \text{if } y_i \in N_C(u) \quad \text{and} \quad x_i \in N_C(b),$$

or

$$y_i \in N_C(v) \quad \text{and} \quad x_i \in N_C(a),$$

and

$$\sigma_i \geq 2 \quad \text{if } y_i \in N_C(v) \quad \text{and} \quad x_i \in N_C(b).$$

Define, for $k \in \{u, v\}$ and $l \in \{a, b\}$,

$$n(kl) := |\{i \in X : y_i \in N_C(k) \text{ and } x_i \in N_C(l)\}|$$

and

$$n(l) := |Y \cap \{i : x_i \in N_C(l)\}|.$$

Then,

$$\begin{aligned} |C| &\geq 2 |N_C(H^*)| + n(ub) + n(va) + 2n(vb) + 2n(a) + 2n(b) \\ &= 2d_C(u) + 2d_C(v) + 2d_C(a) + 2d_C(b) \\ &\quad - (2n(ua) + n(ub) + n(va)). \end{aligned}$$

And $n(ua) + n(ub) \leq d_C(u)$ and $n(ua) + n(va) \leq d_C(a)$. Therefore, $|C| \geq d_C(u) + 2d_C(v) + d_C(a) + 2d_C(b) \geq 6\delta - 6$ and $|G| \geq 6\delta - 2$.

Case 3. $|(y_j, x_j)| = 0$. Then $\sigma_j \geq -1$. Using the argument mentioned in Case 2, one can verify that

$$\begin{aligned} \sigma_i &\geq 1 \quad \text{if } y_i \in N_C(u) \quad \text{and} \quad x_i \in N_C(a), \\ \sigma_i &\geq 2 \quad \text{if } y_i \in N_C(u) \quad \text{and} \quad x_i \in N_C(b), \end{aligned}$$

or

$$y_i \in N_C(v) \quad \text{and} \quad x_i \in N_C(a),$$

and

$$\sigma_i \geq 3 \quad \text{if } y_i \in N_C(v) \quad \text{and} \quad x_i \in N_C(b).$$

Now, one can further compute as in Case 2 to see that $|C| \geq 7\delta - 8 \geq 6\delta - 6$ and $|G| \geq 7\delta - 4 \geq 6\delta - 2$.

5. PROOF OF COROLLARY 1.4

Let C be a longest cycle in the 2-connected triangle-free graph G with $|G| \leq 6\delta - 6$. By Theorem 1.1, it remains to consider the case in which $G - C$ has a unique edge. Let this edge join vertices u and v .

Label $N_C\{u, v\} = \{w_1, w_2, \dots, w_n\}$ in some orientation around C (the subscripts will be reduced modulo n). Clearly, $n \geq 2\delta - 2$ and $|[w_i, w_{i+1}]| \geq 2$ for every i . Divide $N_C\{u, v\}$ into two sets $A := \{w_i : |[w_i, w_{i+1}]| \geq 3\}$ and $B := \{w_i : |[w_i, w_{i+1}]| = 2\}$. Then, $|C| \geq 3|A| + 2|B|$, and $N_H\{w_i, w_{i+1}\}$ is either $\{u\}$ or $\{v\}$ for every $w_i \in B$. If $B = \emptyset$ then $|G| \geq 3n + 2 \geq 6\delta - 4$, a contradiction. Hence, $B \neq \emptyset$. Denote by v_i be the middle vertex of $[w_i, w_{i+1}]$ for each $w_i \in B$.

As in Case 4 of Part II in the proof of Proposition 1.2, one can see that, for any two distinct vertices w_i and w_j of B , v_i and v_j cannot have common vertices in $G - (C \cup H)$. Therefore, if $N_{G-(C \cup H)}(v_i) \neq \emptyset$ for every $i \in B$, then $|G - (C \cup H)| \geq |B|$ and $|G| \geq 3(|A| + |B|) + 2 = 3n + 2 \geq 6\delta - 4$, a contradiction. Hence, $N_{G-(C \cup H)}(v_i) = \emptyset$ for some $i \in B$. Then $[w_{i+1}, w_i]$ and u or v form a longest cycle which is also dominating.

ACKNOWLEDGMENTS

I express my gratitude to Professor H. A. Jung for his many helpful discussions on the problem. Thanks are also due to the referees for their valuable suggestions and comments. Also, I gratefully acknowledge the financial support of the German Academic Exchange Service (DAAD).

REFERENCES

1. P. ASH, Ph. D. thesis, Goldsmiths' College, London, England, 1985.
2. P. ASH AND B. JACKSON, Dominating cycles in bipartite graphs, in "Proc. Silver Jubilee Conference on Combinatorics," University of Waterloo, 1982.
3. J. A. BONDY AND G. FAN, A sufficient condition for dominating cycles, *Discrete Math.* **67** (1987), 205–208.
4. P. FRAISSE, D_λ -cycles and their applications for Hamiltonian graphs, pp. 29–50, D. Sc. thesis, Université De Paris-Sud, 1986.
5. R. HÄGGKVIST, Unsolved problem, in "Proceedings, Fifth Hungarian Colloquium on Combinatorics, 1976."
6. H. A. JUNG, Longest paths joining given vertices in a graph, *Abh. Math. Sem. Univ. Hamburg* **56**, (1986) 127–137.
7. H. A. JUNG, Longest ab -paths in regular graphs, *Graph Theory* (in Memory of G. A. Dirac), in press.
8. H. J. VELDMANN, Existence of dominating cycles and paths in graphs, *Discrete Math.* **43** (1983) 281–296.
9. H. J. VELDMANN, Existence of D_λ -cycles and D_λ -paths, *Discrete Math.* **44** (1983) 309–316.